

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/52, 9/00, 1/21, C12Q 1/68	A3	(11) International Publication Number: WO 00/26381
		(43) International Publication Date: 11 May 2000 (11.05.00)

(21) International Application Number: PCT/US99/25437


(22) International Filing Date: 29 October 1999 (29.10.99)

(30) Priority Data:
60/106,461 30 October 1998 (30.10.98) US(71) Applicant (for all designated States except US): CORNELL
RESEARCH FOUNDATION, INC. [US/US]; Suite 105, 20
Thornwood Drive, Ithaca, NY 14850 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BARANY, Francis
[US/US]; Apartment 12C, 450 E. 63rd Street, New York,
NY 10021 (US). CAO, Weiguo [CN/US]; 420 E. 70th
Street #5N, New York, NY 10021 (US). TONG, Jie
[CN/US]; Apartment 6H, 72-10, 112th Street, Forest Hills,
NY 11375 (US).(74) Agents: GOLDMAN, Michael, L. et al.; Nixon Peabody LLP,
Clinton Square, P.O. Box 1051, Rochester, NY 14603 (US).(81) Designated States: AU, CA, JP, US, European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE).Published
With international search report.(88) Date of publication of the international search report:
9 November 2000 (09.11.00)

(54) Title: HIGH FIDELITY THERMOSTABLE LIGASE AND USES THEREOF



(57) Abstract

The present invention is directed to a thermostable ligase having substantially higher fidelity than either T4 ligase or *Thermus thermophilus* ligase. The DNA molecule encoding this enzyme as well as expression systems and host cells containing it are also disclosed. The thermostable ligase of the present invention is useful in carrying out a ligase detection reaction process and a ligase chain reaction process.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT:

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

NOV 20 2000

HIGH FIDELITY THERMOSTABLE LIGASE AND USES THEREOF

This application claims benefit of U.S. Provisional Patent Application Serial No. 60/106,461 October 30, 1998.

5 The present invention was made with support under National Institutes of Health Grant Nos. GM-41337-09 and PO1-CA65930-02-04). The U.S. Government may have certain rights.

FIELD OF THE INVENTION

10

The present invention is directed to a high fidelity thermostable ligase and uses thereof.

BACKGROUND OF THE INVENTION

15

DNA ligases, as an essential component of DNA replication, recombination, and repair systems found from viruses to humans, catalyze the formation of a phosphodiester bond at single-stranded breaks on duplex DNA (Lehman, I.R., Science, 186:790-797 (1974)). DNA ligases can be classified into two families based on cofactor dependence. ATP-dependent ligases are found in bacteriophages (Dunn, et al., J Mol Biol, 148(4):303-330 (1981) and Weiss, et al., Proc Natl Acad Sci USA, 57(4):1021-1028 (1967)), *Chlorella* virus PBCV-1 (Ho, et al., J Virol, 71(3):1931-19374 (1997)), Vaccinia virus (Shuman, S., Biochemistry, 34(49):16138-161475 (1995)), Archea (Kletzin, A., Nucleic Acids Res, 20(20):5389-5396 (1992) and Bult, et al., Science, 273(5278):1058-1073 (1996)), yeasts (Andaluz, et al., Yeast, 12(9):893-8988 (1996), Ramos, et al., Nucleic Acids Res, 25(8):1485-1492 (1997), Schar, et al., Genes Dev, 11(15):1912-1924 (1997)), mammalian (Tomkinson, et al., Bioessays, 19(10):893-901 (1997), Tomkinson, et al., Mutat Res, 407(1):1-9 (1998), and Wang, et al., J Biol Chem, 269(50):31923-3192811 (1994)), and more recently eubacteria (Cheng, et al., Nucleic Acids Res, 25(7):1369-1374 (1997) and Deckert, et al., Nature, 392(6674):353-358 (1998)). NAD⁺ (i.e. nicotinamide adenine dinucleotide)-dependent ligases, however, are found exclusively

20
25
30

in eubacteria. While some higher eucaryotic organisms may use multiple ATP (i.e. adenosine triphosphate)-dependent ligases to fulfill diverse biological functions, some simple eubacteria genomes could host both an NAD⁺-dependent ligase and an ATP-dependent ligase (Deckert, et al., Nature, 392(6674):353-358 (1998) and Fleischmann, et al., Science, 269(5223):496-512 (1995)). The origin of the additional ATP-dependent ligases in these genomes remains to be determined.

Although the ATP-dependent ligases and NAD⁺-dependent ligases share little sequence homology, all the ligases investigated so far use the same KXDG motif to form adenylated enzyme intermediate (Tomkinson, et al., Bioessays, 19(10):893-901 (1997), Shuman, et al., Virology, 211(1):73-83 (1995), and Luo, et al., Nucleic Acids Res, 24(15):3079-3085 (1996)). Furthermore, they seem to be organized by similar domains and structural folds ((Doherty, et al., Nucleic Acids Res, 24(12):2281-2287 (1996), Subramanya, et al., Cell, 85(4):607-615 (1996), and Sekiguchi, et al., Nucleic Acids Res, 25(4):727-734 (1997)). The diversity of ligase sequences is not only reflected by their different optimal reaction conditions and kinetic rates, but more importantly by their different specificities toward match and mismatch substrates. Among the viral ATP-dependent ligases, the broad substrate tolerance is represented by the T4 enzyme which seals various mismatches on both the 3' and 5' side of the nick junction (Wu, et al., Gene, 76(2):245-254 (1989)).

Vaccinia ligase ligates various mismatches at both 3'-hydroxyl or 5'-phosphate sides with the exception of purine-purine mismatch pairs at the 3'-hydroxyl side (Shuman, S., Biochemistry, 34(49):16138-161475 (1995)). Mammalian ATP-dependent ligases show different substrate sensitivity, as ligase I is more sensitive to 3' mismatches than ligase III (Husain, et al., J Biol Chem, 270(16):9683-9690 (1995)). Additionally, both ligase I and III tolerate a 3'C/T mismatch more than a 3'G/T mismatch. Little is known about archeal ATP-dependent ligases which may reveal the nature of the progenitor of ATP-dependent ligases. Studies on NAD⁺-dependent DNA ligase from *E. coli*, along with T4 ligase, have contributed immensely to understanding of the basic biochemical pathway of the DNA ligation reaction (Lehman, I.R., Science, 186(4166):790-797 (1974) and Rossi, et al., Nucleic Acids Res, 25(11):2106-2113 (1997)). Studies on the NAD⁺-dependent ligase from *Thermus thermophilus* HB8 have revealed the highly discriminative power this enzyme possesses (Luo, et al.,

Nucleic Acids Res, 24(15):3071-3078 (1996)). Although mismatches at 5'-phosphate side are tolerated to some degree (5'A/C, 5'A/A, 5'C/A, 5'C/T, 5'G/T, 5'G/A, 5'T/T, 5'T/G), mismatches at the 3'-hydroxyl side essentially abolish nick-closure activity except 3'G/T or 3'T/G mismatch (Luo, et al., Nucleic Acids Res, 24(15):3071-3078
5 (1996)). Apparently, sequence divergence and subsequent subtle structural variation among DNA ligases underlie an enzyme's recognition preferences toward different mismatched base-pairs.

The study of ligase biochemistry is not only important for understanding its biological functions, but also for developing new technologies. The
10 single nucleotide discrimination observed on DNA ligases has led to the development of ligase-mediated detection techniques (Wu, et al., Gene, 76(2):245-254 (1989), Wu, et al., Genomics, 4(4):560-569 (1989), Landegren, et al., Science, 241(4869):1077-1080 (1988), Landegren, U., Bioessays, 15(11):761-765 (1993), Barany, F., PCR Methods Appl, 1(1):5-16 (1991), and Barany, F., Proc Natl Acad Sci USA, 88(1):189-
15 193 (1991)). Ligase-based linear signal amplification known as LDR (i.e. ligase detection reaction), combined with PCR (i.e. polymerase chain reaction)-based gene specific target amplification, has been proven to be a powerful tool in cancer and disease gene mutation detection (Day, et al., Genomics, 29(1):152-162 (1995)). PCR/LDR technique relies on two properties of a DNA ligase: (i) specificity and (ii)
20 thermostability. *Tth* (i.e. *Thermus thermophilus* HB8) DNA ligase has been successfully used in LDR and LCR (i.e. ligase chain reaction) due to its highly discriminative nick closure activity toward a perfect match substrate and its thermostability which makes thermocycling possible (Barany, F., PCR Methods Appl, 1(1):5-16 (1991) and Barany, F., Proc Natl Acad Sci USA, 88(1):189-193 (1991)).
25 To date, one more ligase was cloned and sequenced from *T. Scot.* (i.e. *Thermus scotoductus*) (Thorbjarnardottir, et al., Gene, 161(1):1-6 (1995) and Jonsson, et al., Gene, 151(1-2):177-180 (1994)), but the substrate specificity of this ligase was not determined.

Despite the existence of a number of ligases from different host
30 sources, the need remains to identify additional ligases with greater fidelity. The present invention is directed to achieving this objective as a result of the cloning and

expression of a ligase from *T. sp.* (i.e. *Thermus* species) AK16D and the biochemical characterization of this high fidelity enzyme.

SUMMARY OF THE INVENTION

5

The present invention is directed to a thermostable ligase having 100 fold higher fidelity than T4 ligase and 6 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent the ligation junction.

10

Another aspect of the present invention is directed to a thermostable ligase having 50 fold higher fidelity than T4 ligase and 5 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base penultimate to the ligation junction.

15

Yet another aspect of the present invention is directed to a thermostable ligase having, in the presence of a Mn^{2+} cofactor, a 12 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent to the ligation junction.

20

The present invention also relates to a DNA molecule encoding the thermostable ligase as well as expression systems and host cells containing such DNA molecules.

25

Another aspect of the present invention relates to the use of the thermostable ligase in carrying out a ligase detection reaction process or a ligase chain reaction process.

30

The ligase detection reaction process, involves detecting a target nucleotide sequence which differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations. This

- 5 -

involves providing a sample potentially containing a target nucleotide sequence which differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations.

- The method further includes providing one or more oligonucleotide probe sets, each characterized by (a) a first oligonucleotide probe having a target specific portion and (b) a second oligonucleotide probe having a target-specific portion. The oligonucleotide probes in a particular set are suitable for hybridization to a target nucleotide sequence which differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations.
- 5 The probes are also suitable for ligation together when hybridized adjacent to one another on the target nucleotide sequence, but have a mismatch which interferes with such ligation when hybridized to any other nucleotide sequence present in the sample.

- The sample, the one or more oligonucleotide probe sets, and the thermostable ligase are blended to form a ligase detection reaction mixture. The ligase detection reaction mixture is subjected to one or more ligase detection reaction cycles comprising a denaturation treatment and a hybridization treatment. In the denaturation treatment, any hybridized oligonucleotides are separated from the target nucleotide sequence. During the hybridization treatment, the oligonucleotide probe sets hybridize at adjacent positions in a base specific manner to their respective target nucleotide sequences, if present in the sample, and ligate to one another. This forms a ligation product sequence containing the target specific portions connected together with the ligation product sequences for each set being distinguishable from other nucleic acids in the ligase detection reaction mixture. The oligonucleotide probe sets may hybridize to a nucleotide sequence in the sample other than their respective target nucleotide sequences but do not ligate together due to a presence of one or more mismatches and individually separate during the denaturation treatment. The presence of ligation product sequences produced as a result of the target nucleotide sequence being present in the sample is then detected.
- 15
20
25

- In the ligase chain reaction process of the present invention, the presence of a target double stranded nucleic acid formed from first and second complementary target nucleotide sequences is detected in a sample. The target double
- 30

stranded nucleic acid differs from other nucleotide sequences by one or more single base changes, insertions, deletions, or translocations.

This method involves providing a sample potentially containing a target double stranded nucleic acid formed from first and second complementary
5 nucleotide sequence. This nucleic acid differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations.

The method further includes providing a first oligonucleotide probe set, characterized by (a) a first oligonucleotide probe having a target specific portion and (b) a second oligonucleotide probe having a target-specific portion. The
10 oligonucleotide probes in the first set are complementary to the first target nucleotide sequence which differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations. The probes are also suitable for ligation together when hybridized adjacent to one another on the first target nucleotide sequence, but have a mismatch which interferes with such ligation
15 when hybridized to any other nucleotide sequence present in the sample. The method of the present invention also requires providing a second oligonucleotide probe set, characterized by (a) a third oligonucleotide probe having a target specific portion and (b) a fourth oligonucleotide probe having a target-specific portion. The oligonucleotide probes in the second set are complementary to the second target
20 nucleotide sequence which differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations. The probes of the second set are suitable for ligation together when hybridized adjacent to one another on the second target nucleotide sequence, but have a mismatch which interferes with such ligation when hybridized to any other nucleotide sequence
25 present in the sample.

The sample, the first and second oligonucleotide probe sets, and the thermostable ligase are blended together to form a ligase chain reaction mixture. The ligase chain reaction mixture is subjected to one or more ligase chain reaction cycles comprising a denaturation treatment and a hybridization treatment. During the
30 denaturation treatment, any hybridized oligonucleotides are separated from the target nucleotide sequences. In the hybridization treatment, the oligonucleotide probe sets hybridize at adjacent positions in a base specific manner to their respective target

nucleotide sequences, if present in the sample. The probes also ligate to one another to form a ligation product sequence containing the target specific portions connected together with the ligation product sequences for each set being distinguishable from other nucleic acids in the ligase chain reaction mixture. The oligonucleotide probe sets may hybridize to nucleotide sequences in the sample other than their respective target nucleotide sequences but do not ligate together due to a presence of one or more mismatches and individually separate during the denaturation treatment. The presence of ligation product sequences produced as a result of the target nucleotide sequence being present in the sample are then detected.

10

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-C show a sequence comparison of *Thermus* DNA ligases. Figure 1A illustrates the evolutionary tree for *Thermus* DNA ligases. Figure 1B is a regional sequence alignment of nine *Thermus* ligases. The aa (i.e. amino acid) sequence of T. scot is retrieved from Genbank by accession number 1085749. The adenylation motif KXDG is underlined and the adenylation site is marked by *. The numbering of aa is based on *Tsp.* AK16D ligase. Figure 1C is a complete amino acid sequence of *Tsp.* AK16D ligase. The adenylation motif KXDG is underlined and the adenylation site ^{118}K is shown with a (*) above the residue. The complete sequence of *Tsp.* AK16D ligase gene and partial sequences of six other *Thermus* ligase genes have been deposited with GenBank under accession No. AF092862 for *Tsp.* AK16D, AF092863 for *Thermus aquaticus* YT-1, AF092864 for *Thermus flavus*, AF092865 for *Thermus filiformis* Tok4A2, AF092866 for *Thermus filiformis* Tok6A1, AF092867 for *Tsp.* Vil3, and AF092868 for *Tsp.* SM32.

Figure 2 shows an SDS-PAGE analysis of *Tsp.* AK16D ligase protein. Lane 1, molecular weight markers; Lane 2, uninduced cell lysate; Lane 3, induced cell lysate; Lane 4, supernatant after heating at 70°C; Lane 5, fraction eluted from Hitrap blue column. The SDS-polyacrylamide gel was 0.1% SDS-7.5% polyacrylamide and was stained with Coomassie brilliant blue after electrophoresis. The arrow points to the location of *Tsp.* AK16D ligase.

30

Figures 3A-C show the effects of salt, pH, and NAD^+ on ligation

activity. *Tsp.* AK16D ligase: closed squares; *Tth* ligase: open squares. Figure 3A reveals the pH effect. Reactions were performed in 20 μ l mixture containing 200 nM nicked duplex substrate, 12.5 pM *Tth* ligase or *Tsp.* AK16D ligase, 20 mM Tris-HCl (pH values were determined at room temperature), 10 mM $MgCl_2$, 100 mM KCl, 10 mM DTT, 1 mM NAD^+ and 20 mg/ml BSA at 65°C for 10 min. Figure 3B shows the salt effect. Reactions were performed in 20 μ l mixture containing 200 nM nicked duplex substrate, 12.5 pM *Tth* ligase or *Tsp.* AK16D ligase, 20 mM Tris-HCl, pH 8.5 (at room temperature) for *Tth* ligase, pH 8.0 for *Tsp.* AK16D ligase, 10 mM $MgCl_2$, indicated amount of KCl, 10 mM DTT, 1 mM NAD^+ and 20 mg/ml BSA at 65°C for 10 min. Figure 3C shows the NAD^+ effect. *Tth* ligation reactions were performed in 20 μ l mixture containing 200 nM nicked duplex substrate, 12.5 pM *Tth* ligase and indicated concentration of NAD^+ , 20 mM Tris-HCl, pH 8.5, 5 mM $MgCl_2$, 100 mM KCl, 10 mM DTT, 1 mM NAD^+ and 20 mg/ml BSA at 65°C for 10 min. *Tsp.* AK16D ligation reaction were performed in 20 μ l mixture containing 200 nM nicked duplex substrate, 12.5 pM *Tth* ligase and indicated concentration of NAD^+ , 20 mM Tris-HCl, pH 8.5, 5 mM $MgCl_2$, 50 mM KCl, 10 mM DTT, 1 mM NAD^+ and 20 mg/ml BSA at 65°C for 10 min.

Figures 4A-B show the divalent cation dependence of *Tsp.* AK16D (stripped bars) and *Tth* (filled bars) ligase activity. Reaction mixtures containing (20 μ l) 20 nM nicked duplex substrate, 0.5 nM *Tth* ligase or 1 nM *Tsp.* AK16D ligase and 5 mM of indicated divalent cation in the reaction buffers as specified in Figure 3C were incubated at 65°C for 10 min. Figure 4A shows the ligation reactions with different divalent ions as the metal cofactor. Figure 4B shows the chromatogram of a representative GeneScan gel illustrating ligation product and DNA adenylate intermediate. (-): negative control reactions in which ligase was omitted. Co^{2+} may have caused precipitation of DNA substrate which resulted in disappearance of the unreacted substrate.

Figures 5A-B shows the time course of *Tth* (Figure 5A) and *Tsp.* AK16D (Figure 5B) ligase activity in the presence of Mg^{2+} (open squares) or Mn^{2+} (closed squares). Reactions were performed in 100 μ l mixture containing 20 nM nicked duplex substrate, 0.5 nM *Tth* ligase or 1 nM *Tsp.* AK16D ligase and 5 mM

Mg²⁺ or Mn²⁺ in the reaction buffers as specified in Figure 3C at 65°C. Aliquots (5 µl) were removed at the indicated time and reactions stopped by adding equal volumes of stop solution.

Figures 6A-B show the divalent cation concentration dependence of *Tth* (Figure 6A) and *Tsp. AK16D* (Figure 6B) ligase activity. Mg²⁺ (open squares); Mn²⁺ (closed squares). Reactions were performed in 20 µl mixture containing 20 nM nicked duplex substrate, 0.5 nM *Tth* ligase or 1 nM *Tsp. AK16D* ligase and indicated concentration of Mg²⁺ or Mn²⁺ in the reaction buffers as specified in Figure 4C at 65°C for 2 min.

Figures 7A-B show the ligation of gapped and inserted substrates. Figure 7A shows the formation of ligated product with gapped and inserted substrates. Reactions were performed in a 20 µl mixture containing 12.5 nM nicked duplex substrate, 1.25 pM *Tth* ligase or 12.5 nM *Tsp. AK16D* ligase in the reaction buffer at 65°C for 4 hours. Figure 7B shows the proposed reaction path leads to ligation of 1 nt (i.e. nucleotides) inserted substrate.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a high fidelity thermostable ligase enzyme. This enzyme has the amino acid sequence of SEQ. ID. No. 1 as follows:

MTLEEARRRVNELRDLIRYHNYLYYVLDAP EISDAEYDRLLRELKELEERFPELKSP
 DSPTEQVGARPLEATFRPVRHPTRMYSLDNAFSLDEVRAFEERIERALGRKGPFlyt
 VERKVDGLSVNLYYEEGILVFGATRGDGETGEEVTQNL LT IPTI PRRLTGVPDRLEV
 RGEVYMPIEAFRLRNQEELEEAGERIFKNPRNAAAGSLRQKDPRVTARRGLRATFYAL
 GLGLEETGLKSQHDLLLWLRERGFVVEHGFTRALGAEGVEEVYQAWLKERRKLPFEA
 DGVVVKLDD LALWRELGYTARTPRFALAYKFPAAEEKETRLLSVAFQVGRTGRITPVG
 VLEPVFIEGSEVSRVTLHNESFIEELDVRIGDWVLVHKAGGV IPEVLRVLKERRTGE
 EKPIIWPENCPECGHALIKEGKVHRCNPPLCPAKRFEAIRHYASRKAMDIQGLGEKL
 IEKLLEKGLVRDVADLYRLKKEDLVNLERMGEKSAENLLRQIEESKGRGLERLLYAL
 GLPGVGEVLARNLALRFGHMDRLLEAGLEDLLEVEGVGELTARAILNTLKDPEFRDL
 VRRLKEAGVEMEAKEREGEALKGLTFVITGELSRPREEVKALLRRLGAKVTDSVSRK
 TSFLVVGENPGSKLEKARALGVPTLSEEELYRLIEERTGKDPRALTA

This protein has a molecular weight of 78 to 81 kDa, as measured by SDS-PAGE. For purposes of the present application, the term "thermostable" refers to a DNA ligase which is resistant to inactivation by heat.

The thermostable ligase of the present invention has a 100 fold higher fidelity than T4 ligase and 6 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent the ligation junction. This ligase also has a 50 fold higher fidelity than T4 ligase and 5 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base penultimate to the ligation junction. Finally, the thermostable ligase of the present invention, in the presence of a Mn^{2+} cofactor, has a 12 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent to the ligation junction. For purposes of the present invention, "fidelity" is defined to mean the ratio of the initial rate of ligating two adjacent probes hybridized to a complementary template with a C-G match at the base of the probe with its 3' end at the ligation junction to the initial rate of ligating two adjacent probes hybridized to a complementary template with a G-T mismatch at the base of the probe with its 3' end at the ligation junction.

The thermostable ligase of the present invention is also characterized by having an arginine adjacent to the active site lysine (i.e. K) in the KXDG motif (where X is any amino acid).

This protein is encoded by a DNA molecule having a nucleotide sequence of SEQ. ID. No. 2 as follows:

ATGACCCTAGAGGAGGCCCGCAGGCGCGTCAACGAACTCAGGGACCTGATCCGTTAC
CACAACCTACCTCTATTACGTCTTGACGCCCGAGATCTCCGACGCCGAGTACGAC
CGGCTCCTTAGGGAGCTTAAGGAGCTGGAGGAGCGCTTCCCGAGCTCAAAGCCCC

GACTCCCCACGGAACAGGTGGGGGCGAGGCCTCTGGAGGCCACCTTCCGCCCGGTG
 CGCCACCCACCCGCATGTACTCCCTGGACAACGCCTTTTCCTTGGACGAGGTGAGG
 GCCTTTGAGGAGCGCATAGAGCGGGCCCTGGGGCGGAAGGGGCCCTTCCTCTACACC
 GTGGAGCGCAAGGTGGACGGTCTTTCGTGAACCTCTACTACGAGGAGGGCATCCTC
 5 GTCTTTGGGGCCACCCGGGGCGACGGGGAGACCGGGGAGGAGGTGACCCAGAACCTC
 CTCACCATCCCCACCATTCCCCGCCGCCTCACGGGCGTTCCGGACCGCCTCGAGGTC
 CGGGGCGAGGTCTACATGCCCATAGAGGCCTTCCTCAGGCTCAACCAGGAGCTGGAG
 GAGGCGGGGAGCGCATCTTCAAAAACCCAGGAACGCCGCCCGCCGGGTCTTGCGG
 CAGAAAGACCCAGGGTCACGGCCAGGCGGGCCCTGAGGGCCACCTTTTACGCCCTG
 10 GGGCTGGGCCTGGAGGAAACCGGGTTAAAAAGCCAGCACGACCTTCTCCTATGGCTA
 AGAGAGCGGGGCTTTCCTGTGAGCACGGCTTTACCGGGGCCCTGGGGGCGGAGGGG
 GTGGAGGAGGTCTACCAGGCCTGGCTCAAGGAGAGGCGGAAGCTTCCCTTTGAGGCC
 GACGGGGTGGTGGTCAAGCTGGACGACCTCGCCCTCTGGCGGGAGCTGGGGTACACC
 GCGCGCACCCCCCGCTTCGCCCTCGCCTACAAGTTCGCCGGCCGAGGAGAAGGAGACC
 15 CGCCTCCTCTCCGTGGCCTTCCAGGTGGGGCGGACCGGGCGCATACCCCCGTGGGC
 GTTCTGGAGCCCGTCTTCATAGAGGGCAGCGAGGTGAGCCGGGTACCCCTCCACAAC
 GAGAGCTTCATTGAGGAGCTGGACGTGCGCATCGGCGACTGGGTGCTGGTCCACAAG
 GCGGGCGGGGTGATTCCCGAGGTGCTGAGGGTCCTGAAAGAGCGCCGACCGGGGAG
 GAGAAGCCCATCATCTGGCCCGAGAACTGCCCGAGTGCGGCCACGCCCTCATCAAG
 20 GAGGGGAAGGTCCACCGCTGCCCAACCCCTTGTGCCCGCCAAGCGCTTTGAGGCC
 ATCCGCCACTACGCCTCCCGCAAGGCCATGGACATCCAGGGCCTGGGGGAGAAGCTC
 ATAGAAAAGCTTCTGGAAGGGCCTGGTCCGGGACGTGGCCGACCTCTACCGCCTG
 AAGAAGGAGGACCTGGTGAACCTGGAGCGCATGGGGGAGAAGAGCGCAGAGAACCTC
 CTCCGCCAGATAGAGGAGAGCAAGGGCCGCGGCCTGGAGCGCCTCCTTTACGCCCTG
 25 GGCCTTCCCGGGGTGGGGGAGGTGCTGGCCCGGAACCTGGCCCTCCGCTTCGGCCAC
 ATGGACCGCCTTCTGGAGGCGGGCCTCGAGGACCTCCTGGAGGTGGAGGGGGTGGGC
 GAGCTCACCGCCCGGGCCATCCTGAATACCCTAAAGGACCCGGAGTTCCGGGACCTG
 GTGCGCCGCCTGAAGGAGGCCGGGGTGGAGATGGAGGCCAAAGAGCGGGAGGGCGAG
 GCCTTGAAGGGGCTCACCTTCGTATCACCGGGGAGCTTTCCCGGCCCGGGAGGAG
 30 GTGAAGGCCCTCCTTAGGCGGCTTGGGGCCAAGGTGACGGACTCGGTGAGCCGCAAG
 ACGAGCTTCTGGTGGTGGGGGAGAACC CGGGGAGCAAGCTGGAAAAGGCCCGCGCC
 TTGGGGGTCCCCACCCTGAGCGAGGAGGAGCTCTACCGCCTCATTGAGGAGAGGACG
 GGCAAGGACCCAAGGGCCCTCACGGCCTAG

35 Fragments of the above polypeptide or protein are also encompassed
 by the present invention.

Suitable fragments can be produced by several means. In the first,
 subclones of the gene encoding the protein of the present invention are produced by
 conventional molecular genetic manipulation by subcloning gene fragments. The
 40 subclones then are expressed *in vitro* or *in vivo* in bacterial cells to yield a smaller
 protein or peptide that can be tested for ligase activity according to the procedure
 described below.

As an alternative, fragments of the ligase of the present invention can
 be produced by digestion of the full-length ligase with proteolytic enzymes like

chymotrypsin or *Staphylococcus* proteinase A, or trypsin. Different proteolytic enzymes are likely to cleave ligase proteins at different sites based on the amino acid sequence of the ligase. Some of the fragments that result from proteolysis may be active ligases.

5 In another approach, based on knowledge of the primary structure of the protein, fragments of the ligase encoding gene may be synthesized by using the PCR technique together with specific sets of primers chosen to represent particular portions of the protein. These then would be cloned into an appropriate vector for increased expression of a truncated peptide or protein.

10 Chemical synthesis can also be used to make suitable fragments. Such a synthesis is carried out using known amino acid sequences for the ligase being produced. Alternatively, subjecting the full length ligase to high temperatures and pressures will produce fragments. These fragments can then be separated by conventional procedures (e.g., chromatography, SDS-PAGE).

15 Variants may also (or alternatively) be modified by, for example, the deletion or addition of amino acids that have minimal influence on the properties, secondary structure and hydrophobic nature of the polypeptide. For example, a polypeptide may be conjugated to a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the
20 protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification, or identification of the polypeptide.

 Suitable DNA molecules are those that hybridize to a DNA molecule comprising a nucleotide sequence of 50 continuous bases of SEQ. ID. No. 2 under stringent conditions characterized by a hybridization buffer comprising 0.9M sodium
25 citrate ("SSC") buffer at a temperature of 37°C and remaining bound when subject to washing with the SSC buffer at 37°C; and preferably in a hybridization buffer comprising 20% formamide in 0.9M saline/0.09M SSC buffer at a temperature of 42°C and remaining bound when subject to washing at 42°C with 0.2x SSC buffer at 42°C.

30 The protein or polypeptide of the present invention is preferably produced in purified form (preferably at least about 80%, more preferably 90%, pure) by conventional techniques. Typically, the protein or polypeptide of the present

invention is secreted into the growth medium of recombinant host cells.

Alternatively, the protein or polypeptide of the present invention is produced but not secreted into growth medium. In such cases, to isolate the protein, the host cell (e.g., *E. coli*) carrying a recombinant plasmid is propagated, lysed by sonication, heat, or chemical treatment, and the homogenate is centrifuged to remove bacterial debris.

The supernatant is then subjected to sequential ammonium sulfate precipitation. The fraction containing the polypeptide or protein of the present invention is subjected to gel filtration in an appropriately sized dextran or polyacrylamide column to separate the proteins. If necessary, the protein fraction may be further purified by HPLC.

10 The DNA molecule encoding the ligase of the present invention can be incorporated in cells using conventional recombinant DNA technology. Generally, this involves inserting the DNA molecule into an expression system to which the DNA molecule is heterologous (i.e. not normally present). The heterologous DNA molecule is inserted into the expression system or vector in proper sense orientation and correct reading frame. The vector contains the necessary elements for the transcription and translation of the inserted protein-coding sequences.

15 U.S. Patent No. 4,237,224 to Cohen and Boyer, which is hereby incorporated by reference, describes the production of expression systems in the form of recombinant plasmids using restriction enzyme cleavage and ligation with DNA ligase. These recombinant plasmids are then introduced by means of transformation and replicated in unicellular cultures including procaryotic organisms and eucaryotic cells grown in tissue culture.

20 Recombinant genes may also be introduced into viruses, such as vaccinia virus. Recombinant viruses can be generated by transfection of plasmids into cells infected with virus.

25 Suitable vectors include, but are not limited to, the following viral vectors such as lambda vector system gt11, gt WES.tB, Charon 4, and plasmid vectors such as pBR322, pBR325, pACYC177, pACYC1084, pUC8, pUC9, pUC18, pUC19, pLG339, pR290, pKC37, pKC101, SV 40, pBluescript II SK +/- or KS +/- (see 30 "Stratagene Cloning Systems" Catalog (1993) from Stratagene, La Jolla, Calif, which is hereby incorporated by reference), pQE, pIH821, pGEX, pET series (see F.W. Studier et. al., "Use of T7 RNA Polymerase to Direct Expression of Cloned Genes,"

Gene Expression Technology vol. 185 (1990), which is hereby incorporated by reference), and any derivatives thereof. Recombinant molecules can be introduced into cells via transformation, particularly transduction, conjugation, mobilization, or electroporation. The DNA sequences are cloned into the vector using standard cloning procedures in the art, as described by Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Springs Laboratory, Cold Springs Harbor, New York (1989), which is hereby incorporated by reference.

A variety of host-vector systems may be utilized to express the protein-encoding sequence(s). Primarily, the vector system must be compatible with the host cell used. Host-vector systems include but are not limited to the following: bacteria transformed with bacteriophage DNA, plasmid DNA, or cosmid DNA; microorganisms such as yeast containing yeast vectors; mammalian cell systems infected with virus (e.g., vaccinia virus, adenovirus, etc.); insect cell systems infected with virus (e.g., baculovirus); and plant cells infected by bacteria. The expression elements of these vectors vary in their strength and specificities. Depending upon the host-vector system utilized, any one of a number of suitable transcription and translation elements can be used.

Different genetic signals and processing events control many levels of gene expression (e.g., DNA transcription and messenger RNA (mRNA) translation).

Transcription of DNA is dependent upon the presence of a promotor which is a DNA sequence that directs the binding of RNA polymerase and thereby promotes mRNA synthesis. The DNA sequences of eucaryotic promotors differ from those of procaryotic promotors. Furthermore, eucaryotic promotors and accompanying genetic signals may not be recognized in or may not function in a procaryotic system, and, further, procaryotic promotors are not recognized and do not function in eucaryotic cells.

Similarly, translation of mRNA in procaryotes depends upon the presence of the proper procaryotic signals which differ from those of eucaryotes. Efficient translation of mRNA in procaryotes requires a ribosome binding site called the Shine-Dalgarno ("SD") sequence on the mRNA. This sequence is a short nucleotide sequence of mRNA that is located before the start codon, usually AUG, which encodes the amino-terminal methionine of the protein. The SD sequences are

complementary to the 3'-end of the 16S rRNA (ribosomal RNA) and probably promote binding of mRNA to ribosomes by duplexing with the rRNA to allow correct positioning of the ribosome. For a review on maximizing gene expression, see Roberts and Lauer, Methods in Enzymology, 68:473 (1979), which is hereby
5 incorporated by reference.

Promoters vary in their "strength" (i.e. their ability to promote transcription). For the purposes of expressing a cloned gene, it is desirable to use strong promoters in order to obtain a high level of transcription and, hence, expression of the gene. Depending upon the host cell system utilized, any one of a
10 number of suitable promoters may be used. For instance, when cloning in *E. coli*, its bacteriophages, or plasmids, promoters such as the T7 phage promoter, *lac* promoter, *trp* promoter, *recA* promoter, ribosomal RNA promoter, the P_R and P_L promoters of coliphage lambda and others, including but not limited, to *lacUV5*, *ompF*, *bla*, *lpp*, and the like, may be used to direct high levels of transcription of adjacent DNA
15 segments. Additionally, a hybrid *trp-lacUV5* (*tac*) promoter or other *E. coli* promoters produced by recombinant DNA or other synthetic DNA techniques may be used to provide for transcription of the inserted gene.

Bacterial host cell strains and expression vectors may be chosen which inhibit the action of the promoter unless specifically induced. In certain operations,
20 the addition of specific inducers is necessary for efficient transcription of the inserted DNA. For example, the *lac* operon is induced by the addition of lactose or IPTG (isopropylthio-beta-D-galactoside). A variety of other operons, such as *trp*, *pro*, etc., are under different controls.

Specific initiation signals are also required for efficient gene
25 transcription and translation in procaryotic cells. These transcription and translation initiation signals may vary in "strength" as measured by the quantity of gene specific messenger RNA and protein synthesized, respectively. The DNA expression vector, which contains a promoter, may also contain any combination of various "strong" transcription and/or translation initiation signals. For instance, efficient translation in
30 *E. coli* requires an SD sequence about 7-9 bases 5' to the initiation codon ("ATG") to provide a ribosome binding site. Thus, any SD-ATG combination that can be utilized by host cell ribosomes may be employed. Such combinations include but are not

...from the L, C, D, E, F, G, H, I, J, K, L, M, N, O, P, Q, R, S, T, U, V, W, X, Y, Z, and other genes. Additionally, any SD
ATG combination produced by recombinant DNA or other techniques involving
incorporation of synthetic nucleotides may be used.

5 Once the isolated DNA molecule encoding the ligase of the present
invention has been cloned into an expression system, it is ready to be incorporated
into a host cell. Such incorporation can be carried out by the various forms of
transformation noted above, depending upon the vector/host cell system. Suitable
host cells include, but are not limited to, bacteria, virus, yeast, mammalian cells,
10 insect, plant, and the like.

 The present invention is useful in a number of processes where a ligase
enzyme is conventionally utilized at high temperatures. Generally, these procedures
include the ligase detection reaction and the ligase chain reaction.

 Both of the ligase detection reaction and ligase chain reaction involve
15 detection of a target sequence and amplification of that sequence at elevated
temperatures. In carrying out these procedures, the enzyme is subjected to elevated
temperatures but is not degraded due to its thermostable character. The ligase
detection reaction and ligase chain reaction procedures are generally described in WO
90/17239 to Barany et. al., F. Barany, et. al., "Cloning, Overexpression, and
20 Nucleotide Sequence of a Thermostable DNA Ligase-Encoding Gene," Gene 109: 1-
11 (1991), and F. Barany, et. al., "Genetic Disease Detection and DNA Amplification
Using Cloned Thermostable Ligase," Proc. Nat'l Acad. Sci. USA 88: 189-93, the
disclosures of which are hereby incorporated by reference.

 The ligase detection reaction process is useful in detecting in a sample
25 a target nucleotide sequence as described more fully below.

 One or more oligonucleotide probe sets are provided for use in
conjunction with this method. Each set includes (a) a first oligonucleotide probe
having a target-specific portion and (b) a second oligonucleotide probe having a
target-specific portion. The oligonucleotide probes in a particular set are suitable for
30 ligation together when hybridized adjacent to one another on a corresponding target
nucleotide sequence, but have a mismatch which interferes with such ligation when
hybridized to any other nucleotide sequence present in the sample.

The sample, the one or more oligonucleotide probe sets, and the ligase are blended to form a ligase detection reaction mixture. The ligase detection reaction mixture is subjected to one or more ligase detection reaction cycles comprising a denaturation treatment and a hybridization treatment. In the denaturation treatment, any hybridized oligonucleotides are separated from the target nucleotide sequences. The hybridization treatment involves hybridizing the oligonucleotide probe sets at adjacent positions in a base-specific manner to the respective target nucleotide sequences, if present in the sample. The hybridized oligonucleotide probes from each set ligate to one another to form a ligation product sequence containing the target-specific portions connected together. The ligation product sequence for each set is distinguishable from other nucleic acids in the ligase detection reaction mixture. The oligonucleotide probe sets may hybridize to adjacent sequences in the sample other than the respective target nucleotide sequences but do not ligate together due to the presence of one or more mismatches. When hybridized oligonucleotide probes do not ligate, they individually separate during the denaturation treatment.

During the ligase detection reaction phase, the denaturation treatment is carried out at a temperature of 80-105°C, while hybridization takes place at 50-85°C. Each cycle comprises a denaturation treatment and a thermal hybridization treatment which in total is from about one to five minutes long. Typically, the ligation detection reaction involves repeatedly denaturing and hybridizing for 2 to 50 cycles. The total time for the ligase detection reaction process is 1 to 250 minutes.

The oligonucleotide probe sets can be in the form of ribonucleotides, deoxynucleotides, modified ribonucleotides, modified deoxyribonucleotides, modified phosphate-sugar-backbone oligonucleotides, nucleotide analogs, and mixtures thereof.

In one variation, the oligonucleotides of the oligonucleotide probe sets each have a hybridization or melting temperature (i.e. T_m) of 66-70°C. These oligonucleotides are 20-28 nucleotides long.

The oligonucleotide probe sets, as noted above, have a reporter label suitable for detection. Useful labels include chromophores, fluorescent moieties, enzymes, antigens, heavy metals, magnetic probes, dyes, phosphorescent groups, radioactive materials, chemiluminescent moieties, and electrochemical detecting moieties.

The product of the ligase detection reaction can be detected in either of two formats. These are fully described in WO 98/03673, to Barany et al., which is hereby incorporated by reference. In one of these formats, ligase detection reaction products are detected by capillary or gel electrophoresis. Alternatively, ligation products can be detected on an array by specific hybridization to a complementary sequence on the array.

The ligation detection reaction mixture may include a carrier DNA, such as salmon sperm DNA.

The hybridization step in the ligase detection reaction, which is preferably a thermal hybridization treatment discriminates between nucleotide sequences based on a distinguishing nucleotide at the ligation junctions. The difference between the target nucleotide sequences can be, for example, a single nucleic acid base difference, a nucleic acid deletion, a nucleic acid insertion, or rearrangement. Such sequence differences involving more than one base can also be detected. Preferably, the oligonucleotide probe sets have substantially the same length so that they hybridize to target nucleotide sequences at substantially similar hybridization conditions. As a result, the process of the present invention is able to detect infectious diseases, genetic diseases, and cancer. It is also useful in environmental monitoring, forensics, and food science.

A wide variety of infectious diseases can be detected by the process of the present invention. Typically, these are caused by bacterial, viral, parasite, and fungal infectious agents. The resistance of various infectious agents to drugs can also be determined using the present invention.

Bacterial infectious agents which can be detected by the present invention include *Escherichia coli*, *Salmonella*, *Shigella*, *Klebsiella*, *Pseudomonas*, *Listeria monocytogenes*, *Mycobacterium tuberculosis*, *Mycobacterium avium-intracellulare*, *Yersinia*, *Francisella*, *Pasteurella*, *Brucella*, *Clostridia*, *Bordetella pertussis*, *Bacteroides*, *Staphylococcus aureus*, *Streptococcus pneumoniae*, B-Hemolytic strep., *Corynebacteria*, *Legionella*, *Mycoplasma*, *Ureaplasma*, *Chlamydia*, *Neisseria gonorrhea*, *Neisseria meningitidis*, *Hemophilus influenza*, *Enterococcus faecalis*, *Proteus vulgaris*, *Proteus mirabilis*, *Helicobacter pylori*, *Treponema*

palladium, *Borrelia burgdorferi*, *Borrelia recurrentis*, *Rickettsial* pathogens, *Nocardia*, and *Acitnomycetes*.

Fungal infectious agents which can be detected by the present invention include *Cryptococcus neoformans*, *Blastomyces dermatitidis*, *Histoplasma capsulatum*, *Coccidioides immitis*, *Paracoccidioides brasiliensis*, *Candida albicans*,
5 *Aspergillus fumigatus*, *Phycomycetes* (*Rhizopus*), *Sporothrix schenckii*, *Chromomycosis*, and *Maduromycosis*.

Viral infectious agents which can be detected by the present invention include human immunodeficiency virus, human T-cell lymphocytotropic virus, hepatitis viruses (e.g., Hepatitis B Virus and Hepatitis C Virus), Epstein-Barr Virus,
10 cytomegalovirus, human papillomaviruses, orthomyxo viruses, paramyxo viruses, adenoviruses, corona viruses, rhabdo viruses, polio viruses, toga viruses, bunya viruses, arena viruses, rubella viruses, and reo viruses.

Parasitic agents which can be detected by the present invention include
15 *Plasmodium falciparum*, *Plasmodium malaria*, *Plasmodium vivax*, *Plasmodium ovale*, *Onchoverva volvulus*, *Leishmania*, *Trypanosoma* spp., *Schistosoma* spp., *Entamoeba histolytica*, *Cryptosporidium*, *Giardia* spp., *Trichimonas* spp., *Balatidium coli*, *Wuchereria bancrofti*, *Toxoplasma* spp., *Enterobius vermicularis*, *Ascaris lumbricoides*, *Trichuris trichiura*, *Dracunculus medinensis*, trematodes,
20 *Diphyllobothrium latum*, *Taenia* spp., *Pneumocystis carinii*, and *Necator americanis*.

The present invention is also useful for detection of drug resistance by infectious agents. For example, vancomycin-resistant *Enterococcus faecium*, methicillin-resistant *Staphylococcus aureus*, penicillin-resistant *Streptococcus pneumoniae*, multi-drug resistant *Mycobacterium tuberculosis*, and AZT-resistant
25 human immunodeficiency virus can all be identified with the present invention.

Genetic diseases can also be detected by the process of the present invention. This can be carried out by prenatal or post-natal screening for chromosomal and genetic aberrations or for genetic diseases. Examples of detectable genetic diseases include: 21 hydroxylase deficiency, cystic fibrosis, Fragile X
30 Syndrome, Turner Syndrome, Duchenne Muscular Dystrophy, Down Syndrome or other trisomies, heart disease, single gene diseases, HLA typing, phenylketonuria, sickle cell anemia, Tay-Sachs Disease, thalassemia, Klinefelter Syndrome,

Huntington Disease, autoimmune diseases, lipidosis, obesity defects, hemophilia, inborn errors of metabolism, and diabetes.

Cancers which can be detected by the process of the present invention generally involve oncogenes, tumor suppressor genes, or genes involved in DNA
5 amplification, replication, recombination, or repair. Examples of these include: BRCA1 gene, p53 gene, APC gene, Her2/Neu amplification, Bcr/Ab1, K-ras gene, and human papillomavirus Types 16 and 18. Various aspects of the present invention can be used to identify amplifications, large deletions as well as point mutations and small deletions/insertions of the above genes in the following common human
10 cancers: leukemia, colon cancer, breast cancer, lung cancer, prostate cancer, brain tumors, central nervous system tumors, bladder tumors, melanomas, liver cancer, osteosarcoma and other bone cancers, testicular and ovarian carcinomas, head and neck tumors, and cervical neoplasms.

In the area of environmental monitoring, the present invention can be
15 used for detection, identification, and monitoring of pathogenic and indigenous microorganisms in natural and engineered ecosystems and microcosms such as in municipal waste water purification systems and water reservoirs or in polluted areas undergoing bioremediation. It is also possible to detect plasmids containing genes that can metabolize xenobiotics, to monitor specific target microorganisms in
20 population dynamic studies, or either to detect, identify, or monitor genetically modified microorganisms in the environment and in industrial plants.

The present invention can also be used in a variety of forensic areas, including for human identification for military personnel and criminal investigation, paternity testing and family relation analysis, HLA compatibility typing, and
25 screening blood, sperm, or transplantation organs for contamination.

In the food and feed industry, the present invention has a wide variety of applications. For example, it can be used for identification and characterization of production organisms such as yeast for production of beer, wine, cheese, yogurt, bread, etc. Another area of use is with regard to quality control and certification of
30 products and processes (e.g., livestock, pasteurization, and meat processing) for contaminants. Other uses include the characterization of plants, bulbs, and seeds for

breeding purposes, identification of the presence of plant-specific pathogens, and detection and identification of veterinary infections.

Desirably, the oligonucleotide probes are suitable for ligation together at a ligation junction when hybridized adjacent to one another on a corresponding target nucleotide sequence due to perfect complementarity at the ligation junction. However, when the oligonucleotide probes in the set are hybridized to any other nucleotide sequence present in the sample, there is a mismatch at a base at the ligation junction which interferes with ligation. Most preferably, the mismatch is at the base at the 3' base at the ligation junction. Alternatively, the mismatch can be at the bases adjacent to bases at the ligation junction.

Before carrying out the ligase detection reaction, in accordance with the present invention, target nucleotide sequences in the sample can be preliminarily amplified. This preferably carried out with polymerase chain reaction. The polymerase chain reaction process is fully described in H. Erlich, et. al, "Recent Advances in the Polymerase Chain Reaction," Science 252: 1643-50 (1991); M. Innis, et. al., PCR Protocols: A Guide to Methods and Applications, Academic Press: New York (1990) and R. Saiki, et. al., "Primer-directed Enzymatic Amplification of DNA with a Thermostable DNA Polymerase," Science 239: 487-91 (1988), which are hereby incorporated by reference.

The ligase detection reaction process achieves linear amplification of a target sequence. The ligase chain reaction utilizes essentially the same steps and conditions as the ligase detection reaction to achieve exponential amplification. This greater level of amplification is achieved by utilizing 2 sets of complementary oligonucleotides with each set hybridizing to complementary strands of a target nucleic acid sequence.

In the ligase chain reaction process of the present invention, the presence of a target double stranded nucleic acid formed from first and second complementary target nucleotide sequences is detected in a sample. The target double stranded nucleic acid differs from other nucleotide sequences by one or more single base changes, insertions, deletions, or translocations.

This method involves providing a sample potentially containing a target double stranded nucleic acid formed from first and second complementary

nucleotide sequence. This nucleic acid differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations.

The method further includes providing a first oligonucleotide probe set, characterized by (a) a first oligonucleotide probe having a target specific portion and (b) a second oligonucleotide probe having a target-specific portion. The oligonucleotide probes in the first set are complementary to the first target nucleotide sequence which differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations. The probes are also suitable for ligation together when hybridized adjacent to one another on the first target nucleotide sequence, but have a mismatch which interferes with such ligation when hybridized to any other nucleotide sequence present in the sample. The method of the present invention also requires providing a second oligonucleotide probe-set, characterized by (a) a third oligonucleotide probe having a target specific portion and (b) a fourth oligonucleotide probe having a target-specific portion. The oligonucleotide probes in the second set are complementary to the second target nucleotide sequence which differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations. The probes of the second set are suitable for ligation together when hybridized adjacent to one another on the second target nucleotide sequence, but have a mismatch which interferes with such ligation when hybridized to any other nucleotide sequence present in the sample.

The sample, the first and second oligonucleotide probe sets, and the thermostable ligase are blended together to form a ligase chain reaction mixture. The ligase chain reaction mixture is subjected to one or more ligase chain reaction cycles comprising a denaturation treatment and a hybridization treatment. During the denaturation treatment, any hybridized oligonucleotides are separated from the target nucleotide sequences. In the hybridization treatment, the oligonucleotide probe sets hybridize at adjacent positions in a base specific manner to their respective target nucleotide sequences, if present in the sample. The probes also ligate to one another to form a ligation product sequence containing the target specific portions connected together with the ligation product sequences for each set being distinguishable from other nucleic acids in the ligase chain reaction mixture. The oligonucleotide probe

sets may hybridize to nucleotide sequences in the sample other than their respective target nucleotide sequences but do not ligate together due to a presence of one or more mismatches and individually separate during the denaturation treatment. The presence of ligation product sequences produced as a result of the target nucleotide sequence being present in the sample are then detected.

EXAMPLES

Example 1 - Reagents, Media, and Strains

All routine chemical reagents were purchased from Sigma Chemicals (St. Louis, MO) or Fisher Scientific (Fair Lawn, NJ). Restriction enzymes and T4 DNA ligase were obtained from New England Biolabs (Beverly, MA). Oligonucleotide synthesis reagents, DNA sequencing kits, and PCR kits were obtained from Applied Biosystems Division of Perkin-Elmer Corporation (Foster City, CA). dNTPs, BSA (i.e. bovine serum albumin), ATP were purchased from Boehringer-Mannheim (Indianapolis, IN). *Pfu* DNA polymerase was purchased from Stratagene (La Jolla, CA). *E. coli* strain NovaBlue(DE3)pLysS, and plasmid pET11c were purchased from Novagen, Inc. (Madison, WI). Protein assay kit was from Bio-Rad (Hercules, CA). HiTrap Blue affinity column was from Pharmacia (Piscataway, NJ). LB medium was prepared according to standard formula (Sambrook, et al., (1989) Molecular Cloning-A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1994), which is hereby incorporated by reference). Sonication buffer consisted of 50 mM Tris-HCl, pH 8.0 and 1 mM EDTA. TE buffer consisted of 10 mM Tris-HCl, pH 8.0 and 1 mM EDTA. *Tth* DNA ligase and its mutant K294R were purified as previously described (Luo, et al., Nucleic Acids Res., 24(15):3071-3078 (1996), which is hereby incorporated by reference).

Example 2 - Oligonucleotide Synthesis

Oligonucleotides were synthesized by using a 394 automated DNA synthesizer from Applied Biosystems Division of Perkin-Elmer Corp. PCR and sequencing primers were purified by ethanol precipitation according to instruction

manual. The degenerate sense primer 5'-ATC(T/A)(C/G)CGACGC(C/G)-
GA(G/A)TA(T/C)GA-3' (SEQ. ID. No. 3) corresponding to amino acids 32-38
(ISDAEYD) (SEQ. ID. No. 4) in the *T. thermophilus* HB8 DNA ligase gene, and
antisense primers 5'-CC(C/G)GT(C/G)C(G/T)-(G/C)CC(G/C)AC(C/T)TG(A/G)AA-
5 3' (SEQ. ID. No. 5) and 5'-GCCTTCTC(C/G/A)A(A/G)(T/C)TTG-
(C/G)(A/T)(G/C)CC-3' (SEQ. ID. No. 6) corresponding to amino acids 333-339
(FQVGRTG) (SEQ. ID. No. 7) and 641-647 (GSKLEKA) (SEQ. ID. No. 8) were
used to amplify DNA ligase gene fragments from *Thermus* strains. Additional PCR
and sequencing primers were synthesized as required. PCR amplification primers for
10 cloning *Tsp.* AK16D DNA ligase gene into pET11c vector were 5'-
GCGATTTCATATGACCCTAGAGGAGGCCCG-3' (SEQ. ID. No. 9) and 5'-
GCGGGATCCGAGGC CTTGGAGAAGCTCTT-3', (SEQ. ID. No. 10) where the
NdeI and *BamHI* sites are underlined and the initiation codon in the forward primer is
shown in bold. Oligonucleotide substrates for ligation assay were purified on a
15 denaturing sequencing gel (7 M urea/10% polyacrylamide) (Applied Biosystems Inc.,
The complete guide to evaluating and isolating synthetic oligonucleotides, Applied
Biosystems Inc., Foster City, CA (1992)). 5'-phosphorylation of oligonucleotides was
achieved during synthesis by using Chemical Phosphorylation Reagent (Glen
Research, Sterling, VA). Fluorescent group was attached to a 3'-terminus using
20 Fluorescein CPG column (Glen Research).

Example 3 - DNA Amplification, Cloning And Sequence Analysis

Genomic DNAs from *Thermus* strains were isolated as previously
25 described (Cao, et al., Gene, 197:205-214 (1997), which is hereby incorporated by
reference). PCR amplifications with degenerate and unique primers and inverse PCR
on circularized templates were carried out in a GeneAmp PCR System 9700
thermocycler (Applied Biosystems Division of Perkin Elmer) as described (Wetmur,
et al., J Biol Chem, 269(41):25928-25935 (1994), which is hereby incorporated by
30 reference). The nucleotide sequences of amplified ligase fragments were directly
determined on an ABI 373 sequencer using ABI PRISM™ Dye Terminator Cycle
Sequencing Ready Reaction Kit (Perkin-Elmer). Full length *Tsp.* AK16D DNA
ligase gene was amplified using PCR amplification primers as described above,

- 25 -

digested with *Nde*I and *Bam*HI, ligated into the cloning vector pET11c treated with the same pair of restriction enzymes, and transformed into *E. coli* strain NovaBlue(DE3)pLysS. Inserts in pET expression vectors were sequenced in both orientations to ensure that the plasmid constructs were free of PCR or ligation error.

5 Nucleic acid and protein sequence analyses were carried out by Clustal method (Higgins, et al., Comput Appl Biosci, 5(2):151-153 (1989), which is hereby incorporated by reference) using MegAlign program of DNASTAR (Madison, WI).

Example 4 - Expression and Purification of *Tsp.* AK16D DNA Ligase

10 *E. coli* NovaBlue(DE3)pLysS cells containing plasmid pTAK encoding the *Tsp.* AK16D DNA ligase gene from a pET11c construct was propagated overnight at 37°C in LB medium containing 50 µg/ml ampicillin, 25 µg/ml chloramphenicol, and 0.2% glucose. Overnight cultures were diluted 100-fold into

15 the same medium, grown until the optical density of the culture reached 0.5 at 600 nm, then induced by the addition of IPTG to a final concentration of 1 mM, and grown for an additional 4 hrs under the same conditions. Cells were collected by centrifugation, frozen/thawed at -20°C/23°C, disrupted by sonication, and clarified by centrifugation as previously described (Wetmur, et al., J Biol Chem, 269(41):25928-

20 25935 (1994), which is hereby incorporated by reference). The resulting supernatants were heated at 70°C for 15 min to denature thermolabile *E. coli* proteins, placed on ice for 30 min to aggregate the denatured proteins, and cleared of denatured proteins by microcentrifugation for 15 min at 4°C. The partially pure DNA ligase was further purified by chromatography using 1 ml HiTrap Blue affinity column. Briefly, the

25 column containing *Tsp.* AK16D DNA ligase was washed extensively with TE buffer (pH 7.8) containing 0.1 M NaOAc, and the ligase was eluted with TE buffer (pH 7.8) containing 2 M NaCl. After dialysis against TE buffer (pH 8.0) containing 0.2 M KCl and concentration using Centricon-30 (Amicon), protein concentration was assayed by the Bradford method with reagents supplied by Bio-Rad protein assay kit. The

30 amount of protein was determined using BSA as the standard. The purity of the ligase was verified through 7.5% SDS (i.e. sodium dodecyl sulfate)-PAGE (i.e. polyacrylamide gel electrophoresis) analysis followed by visualizing the overloaded gel with routine Coomassie Brilliant Blue R staining.

Example 5 - Substrates And Ligation Assay

The oligonucleotide perfect match substrate was formed by annealing two short oligonucleotides (33-mer for LP3'C (SEQ. ID. No. 11) and 30-mer for Com3F (SEQ. ID. No. 12)) with a 59-mer complementary oligonucleotide (Glg). Oligonucleotides LP3'C and Glg (SEQ. ID. No. 14) were in 1.5-fold excess so that the all the 3' Fam labeled Com3F represented nicked substrates (see Luo, et al., Nucleic Acids Res, 24(15):3071-3078 (1996), which is hereby incorporated by reference).

The T/G mismatch substrate was formed by annealing LP3'T (SEQ. ID. No. 13), which introduced a single base-pair mismatch at the 3'-end of the nick junction, along with Com 3'F to the complementary strand (Glg). The nicked DNA duplex substrates were formed by denaturing DNA probes at 94°C for 2 min followed by re-annealing at 65°C for 2 min in ligation buffer. The sequences of the oligonucleotides were listed below (p represents 5' phosphate group):

		PAGTTGTCATAGTTTGATCCTCTAGTCTGGG-Fam-3' Com3
LP3'T	5'-CCCTGTTCCAGCGTCTGCGGTGTTGCGTT	
LP3'C	5'-AAAACCTGTTCCAGCGTCTGCGGTGTTGCGTC	
Glg	3'-GGGACAAGGTCGCAGACGCCACAACGCAGTCAACAGTATCAAAGTAGGAGATCAGACCC-5'	

Ligation mixtures (20 µl) containing indicated amount of DNA ligase and match or mismatch substrate in the ligase buffer (20 mM Tris-HCl, pH 7.6 at room temperature; 10 mM MgCl₂; 100 mM KCl; 10 mM DTT (i.e. dithiothreitol); 1 mM NAD⁺; and 20 mg/ml BSA) were incubated at 65°C for a predetermined time. Reactions were terminated by the addition of an equal volume of stop solution (i.e. 50 mM EDTA, 80% formamide, and 1% Blue Dextran). Samples (5 µl) were electrophoresed through an 8 M urea-10% polyacrylamide GeneScan gel according to instructional manual (Perkin Elmer). The unreacted substrates were represented by the 30-mer com3F and products were represented by a ligated 63-mer in the case of the match substrate. Both the remaining substrates and ligated products were quantified using GeneScan analysis software 672 (version 2.0, Perkin Elmer).

Example 6 - Steady State Kinetics

Steady state kinetic constants were determined by measuring initial rates of the ligation reaction at a given substrate concentration (nicked DNA duplex substrate concentration ranging from 25-400 nM) and a given ligase concentration (12.5 pM for both *Tth* and *Tsp*. AK16D) in 100 µl reaction volume at 65°C. A 5 µl aliquot was removed at 0, 2, 4, 6, 8, 10 min, and mixed with 5µl of stop solution. The remaining substrate was separated from ligated product by GeneScan gel as described above. Initial rates of the ligation reactions were calculated from the generation of ligated product over time. The K_m and k_{cat} values were determined using computer software Ultrafit (Biosoft, Ferguson, MO).

Example 7 - Sequence Analysis Of Seven *Thermus* Ligase Genes

Amino acid sequence alignment of five Gram negative bacterial NAD⁺-dependent DNA ligases indicates that *Tth* ligase is 93% identical to *Thermus scotoductus* ligase, 49% to *Rhodothermus marinus* ligase, 48% to *E. coli* ligase, and 38% to *Zymomonas mobilis* based on sequence data retrieved from GeneBank. Degenerate primers corresponding to highly conserved regions of these ligases were used to amplify fragments of ligase genes from seven *Thermus* strains which represent a worldwide collection: *Thermus flavus* from Japan (SEQ. ID. No. 16), *Thermus aquaticus* YT-1 (SEQ. ID. No. 15) and *Thermus sp.* AK16D from Yellowstone National Park in the United States, *Thermus filiformis* Tok4A2 (SEQ. ID. No. 17) and *Thermus filiformis* Tok6A1 (SEQ. ID. No. 18) from New Zealand, *Thermus sp.* SM32 (SEQ. ID. No. 19) from Azores, and *Thermus sp.* Vil3 (SEQ. ID. No. 20) from Portugal. The sequences of amplified ligase fragments ranging from 1.4 to 1.6 kb were determined by directly sequencing the PCR products using an ABI 373 automated sequencer. *Thermus* ligases, in general, were highly conserved during evolution as demonstrated by 85%-98% sequence identity. In contrast, the amino acid sequences of the restriction endonuclease *TaqI* and its isoschizomers from the identical strains show only 50-70% aa identities (Cao, et al., *Gene*, 197:205-214 (1997), which is hereby incorporated by reference). *Thermus* ligases in general show 30-40% sequence identities as compared with DNA ligases from other bacteria. The

sequence divergence is slightly higher among the different geographic groups than within the same group, which may reflect random drift or adaptation to their respective local environments (Figure 1). *Thermus flavus*, *Thermus filiformis* Tok4A2, *Thermus filiformis* Tok6A1, *Thermus sp.* SM32, *Thermus sp.* Vil3, *Thermus aquaticus* YT-1, and *Thermus sp.* AK16D (SEQ. ID. No. 14) ligases shared 98.2%, 89.9%, 89.5%, 89.8%, 88.3%, 88.2%, 88.1% with *Thermus thermophilus* HB8 DNA ligase, respectively. The adenylation site of the enzymes (¹¹⁸KXDG where X is in general a hydrophobic residue), as identified by site-directed mutagenesis of *Tth* DNA ligase, is completely identical among all *Thermus* ligases, furthermore, the flanking sequences of the adenylation motif are also identical except *Tsp.* AK16D in which the aa residues ¹¹⁷H before the ¹¹⁸K is substituted by an ¹¹⁸R (Figure 1B). In non-*Thermus* NAD⁺-dependent ligases discovered to date, the corresponding position is either a Pro or a Leu. The two isolates from Japan can be distinguished from the other *Thermus* strains by a 3-aa-insertion at position 234.

15

Example 8 - Cloning, Expression And Purification Of DNA Ligase From *Tsp.* AK16D

To maximize the chance of finding a *Thermus* ligase with novel properties, *Tsp.* AK16D ligase was chosen which showed the least sequence identity as compared with *T. thermophilus* ligase. To obtain the complete sequence of the ORF (i.e. open reading frame), the fragments of the N- and C-terminus of the gene were amplified by inverse PCR and were subject to direct sequencing. The complete ORF of the *Thermus sp.* AK16D ligase gene consists of 674 amino acids, as compared to 676 aa for *Tth* ligase and 674 aa for *T. scot* ligase (Figure 1C). The full-length *Thermus sp.* AK16D ligase gene was PCR amplified using *Pfu* polymerase and cloned into expression plasmid pET11c (Novagen). The integrity of the insert containing the ligase gene was verified by DNA sequencing. The pET11c plasmid expressing *Tsp.* AK16D ligase was transformed into competent *E. coli* cells NovaBlue(DE3)pLysS. Production of ligases was induced by adding IPTG to 1 mM final concentration. *Tsp.* AK16D ligase protein was expressed to approximately 10% of total cellular proteins (Figure 2, lane 3). Heating at 70°C for 15 minutes denatured most of *E. coli* proteins while leaving the thermostable ligases as the dominant band (Figure 2, lane 4). A

cibacron blue based affinity chromatography (Pharmacia) further removed residual *E. coli* proteins and nucleic acids, yielding apparently homogenous *Tsp. AK16D* ligase protein as judged by Coomassie staining (Figure 2, lane 5).

5 **Example 9 - Salt, pH, and NAD⁺ Dependence Of The Ligation Reaction**

Figure 3A depicts the pH dependence of ligase activity of *Tth* and *Tsp. AK16D* ligase proteins. The shape of the pH dependence curves of *Tth* ligase and *Tsp. AK16D* ligase is essentially superimposable. The optimal pH is 8.5 for both *Tth*
10 ligase and *Tsp. AK16D* ligase with greater than 80% activity observed between pH 7.8 and 9.5. The identity of pH effect suggests that both of the ligases possess similar local environment at their catalytic center, which is in agreement with the degree of sequence conservation between the two ligases. Figure 3B depicts the salt concentration dependence of ligase activity of *Tth* and *Tsp. AK16D* ligase proteins.
15 The optimum KCl concentration for *Tth* ligase and *Tsp. AK16D* ligase are 100 and 50 mM, respectively. Figure 3C depicts the NAD⁺ concentration dependence of ligase activity of *Tth* and *Tsp. AK16D* proteins. The optimum NAD⁺ concentration is 1 mM for both *Tth* ligase and *Tsp. AK16D* ligase. The similarity of the NAD profiles is in keeping with the highly conserved nature of the N-terminal domain of the ligases
20 which is involved in NAD⁺ binding.

Example 10 - Effects Of Divalent Metals On The Ligation Reaction

Divalent metal ion is indispensable for each of the three steps in a
25 ligation reaction: (i) adenylation of a lysine residue in the adenylation motif KXDG; (ii) transfer of the adenylate to the 5' phosphate to form a DNA-adenylate intermediate; and (iii) formation of the phosphodiester bond with the release of adenosine monophosphate (AMP). In general, Mg²⁺ is the preferred metal ion for both ATP-dependent and NAD⁺-dependent ligases. Mg²⁺ was substituted with
30 alkaline earth metal ion Ca²⁺ and commonly studied period 4 transition metal ions. *Tth* and *Tsp. AK16D* ligases could use Mn²⁺ as an alternative metal cofactor to support ligation activity (Figure 4). Both enzymes were less active with Ca²⁺, while Co²⁺, Ni²⁺, Cu²⁺, and Zn²⁺ failed to support ligation. In comparison, ATP-dependent

ligase from Hin (i.e. *Haemophilus influenzae*) uses only Mg^{2+} and Mn^{2+} as the metal cofactor for nick closure but not Ca^{2+} , Co^{2+} , Cu^{2+} , and Zn^{2+} (Cheng, et al., Nucleic Acids Res, 25(7):1369-1374 (1997), which is hereby incorporated by reference); ATP-dependent ligase from *Chlorella* virus PBCV-1 can use Mg^{2+} , Mn^{2+} , and Co^{2+} but not Ca^{2+} , Cu^{2+} , and Zn^{2+} (Ho, et al., J Virol, 71(3):1931-1937 (1997), which is hereby incorporated by reference). Using Ca^{2+} as the metal cofactor, *Thermus* enzymes were able to convert most of the substrate into the DNA-adenylate intermediate. However, the rates of nick closure were reduced which led to the accumulation of the DNA-adenylate intermediate (Figure 4B). A small amount of the intermediate was observed with Ni^{2+} ; however, ligation product was not observed at the current detection level, suggesting that Ni^{2+} could not support the nick closure step (Figure 4B). To further compare the relative activity of the two *Thermus* ligases with Mg^{2+} and Mn^{2+} , the generation of ligation product was first monitored over a 20-min time period. As shown in Figure 5, the *Thermus* enzymes were consistently more active with Mg^{2+} than with Mn^{2+} . Second, ligation activity up to 40 mM Mg^{2+} or Mn^{2+} concentrations (Figure 6) was assayed. Both of the enzymes responded sensitively to the change of the metal ion concentration in the reaction mixture. At high M^{2+} concentrations, the high ionic strength may inhibit the enzyme activity, consistent with KCl dependence profile (Figure 4). Similar to the time-course results, the *Thermus* enzymes were more active with Mg^{2+} than with Mn^{2+} (Figure 6). The discrepancy on the relative activity of *Thermus* ligases between this study and an earlier report may be due to use here of cloned enzymes while the earlier work used purified native enzyme (Takahashi, et al., J Biol Chem, 259(16):10041-10047 (1984), which is hereby incorporated by reference).

Example 11 - Steady State Kinetics

The steady state kinetic constants were measured by monitoring the formation of fluorescently labeled ligation product over time using substrate concentrations spanning estimated K_m values (Table 1).

Table 1. Steady state kinetics of *Tth* and *Tsp*. AK16D ligase ^a

Ligase	K_m (nM)	k_{cat} (min ⁻¹)	k_{cat}/K_m (M ⁻¹ .s ⁻¹)
<i>Tth</i>	87	56	1.1×10^7
<i>Tsp</i> .AK16D	104	38	0.62×10^7

^a: Results represent the average of at least three experiments.

5 The steady state properties of *Tsp*. AK16D ligase were similar to *Tth* ligase, indicating that the catalytic channels are highly conserved in *Thermus* ligases. The average K_m value of about 90 nM for *Thermus* ligases is similar to the K_m value of 50 nM for *E. coli* ligase (Modrich, et al., J Biol Chem, 248(21):7495-7501 (1973), which is hereby incorporated by reference) and about 10-fold higher than vaccinia virus ATP-dependent ligase (Sekiguchi, et al., Nucleic Acids Res, 25(4):727-734 (1997), which is hereby incorporated by reference). The average k_{cat} value of about 45 turnovers per min for *Thermus* ligases is higher than the k_{cat} value of 28 turnovers per min for *E. coli* ligase (Modrich, et al., J Biol Chem, 248(21):7495-7501 (1973), which is hereby incorporated by reference).

Example 12 - Ligation Of Gapped Or Inserted DNA Duplex Substrates

15 Gapped substrates were formed by deleting one or two nt from the 3' hydroxyl site of oligonucleotide LP3'C, and inserted substrates were formed by adding one or two nt at the 3' hydroxyl site of oligonucleotide LP3'C. Gapped or inserted duplexed DNA sequences are distinctively different from normal nicked substrate. Under our experimental conditions, no ligation was detectable with 1-nt (i.e. nucleotide) or 2-nt gapped or 2-nt insertion substrates for either *Tth* or *Tsp*. AK16D ligase (Figure 7A). As for 1-nt insertion substrates, only A insertion gave a trace amount of ligated products for both ligases (Figure 7A). All other 1-nt insertions at the ligation junction could not be ligated. In contrast, *Hin* ligase and *Chlorella* ligase demonstrate observable ligation with 1-nt gap (Ho, et al., J Virol, 71(3):1931-1937 (1997) and Cheng, et al., Nucleic Acids Res, 25(7):1369-1374 (1997), which are hereby incorporated by reference). In the case of vaccinia ligase, the ligation of 1-nt gap is negligible but the formation of DNA-adenylate intermediate is significant, suggesting the major impact of using 1-nt gapped substrate is on nick

closure (Shuman, S., Biochemistry, 34(49):16138-16147 (1995), which is hereby incorporated by reference). The formation of DNA-adenylate intermediate with the *Thermus* enzymes was not observed, suggesting that most of the gapped or inserted substrates may have abolished the possibility of completing the second step in the ligation cycle — adenylation of DNA substrate at the 5' phosphate. The 1-nt A insertion mis-ligation could be due to slippage (Figure 7B). Although *Thermus* ligase slippage is far less than *Thermus* DNA polymerase, it does occur at a low frequency. Given the fact that the adjacent nt is a T, the slippage could have occurred at 5' phosphate side where a 5'A/C mismatch is ligated (Luo, et al., Nucleic Acids Res, 24(15):3071-3078 (1996), which is hereby incorporated by reference). It is unlikely that the enzyme tolerates slippage on the 3' side, because a 1 nt C insertion did not yield detectable ligation product (Figure 7).


Example 13 - *Thermus* DNA Ligase Fidelity

Tth DNA ligase is more discriminative when the mismatch is located at the 3' side of the nick junction. 3'G/T or 3'T/G is the only mismatch that shows observable mismatch ligation (Luo, et al., Nucleic Acids Res, 24(15):3071-3078 (1996), which is hereby incorporated by reference). To evaluate the fidelity of the cloned *Tsp.* AK16D ligase, the rate ratio of match over 3'T/G mismatch ligation was compared with wild-type and K294R mutant *Tth* DNA ligases along with T4 ligase from a commercial source (Table 2).

Table 2. DNA ligase fidelity ^a

Ligase	Enzyme Concentration (nM)	Initial rates of C-G match (fmol/min)	Initial rates of T-G mismatch at 3'-end (fmol/min)	Initial rates of T-G mismatch at penultimate 3'-end (fmol/min)	Ligation fidelity 1 ^b	Ligation fidelity 2 ^c
T4	0.5	1.4×10^2	2.8	7.1	5.0×10^1	1.9×10^1
<i>Tth</i> -wt	1.25	5.5×10^1	6.5×10^{-2}	2.9×10^{-1}	8.4×10^2	1.9×10^2
<i>Tth</i> -K294R	12.5	1.5×10^2	2.3×10^{-2}	4.3×10^{-1}	6.3×10^3	3.4×10^2
<i>Tsp</i> . AK16D	12.5	1.3×10^2	2.5×10^{-2}	1.2×10^{-1}	5.1×10^3	1.1×10^3

^a The reaction mixture consisted of 12.5 nM nicked DNA duplex substrates, indicated the amount of DNA ligases in ligation reaction buffer. T4 DNA ligase fidelity was assayed at 37 °C, thermophilic ligase fidelity was assayed at 65°C. Five μ l Aliquots from a 160 μ l reaction mixture were removed at 0, 10, 20, 30, 40, 50, 60 s for reactions containing matched substrates and at 0, 1, 2, 3, 4, 5, 6 h for reactions containing mismatched substrates, and mixed with 5 μ l of stop solution. Samples (5 μ l) were electrophoresed through an 8 M urea-10% polyacrylamide gel as described. Fluorescently labeled ligation products were analyzed and quantified using Genescan 672 version 2.0 software (Applied Biosystems, Foster City, CA). The results were plotted using DeltaGraph Pro3 software (DeltaPoint Inc., Monterey, CA). The initial rates were determined as the slope of linear range of the graph with the x-axis as the time and the y-axis as the amount of the ligation product generated. A schematic illustration of matched and mismatched substrates are as follows:



^b Ligation fidelity 1= Initial Rate of C-G match / Initial Rate of T-G mismatch at 3'-end.

^c Ligation fidelity 2= Initial Rate of C-G match / Initial Rate of T-G mismatch at penultimate 3'-end. The concentrations of DNA ligases used in each experiment are as indicated. Results were calculated as the average of at least two experiments.

T4 ligase demonstrated high catalytic efficiency toward both match and 3'T/G mismatch substrate such that a ligation fidelity of 50 was obtained. *Thermus* ligases appeared to be less efficient in match ligation as evidenced by the requirement of higher enzyme concentration to achieve comparable match ligation rate. However, under the same assay conditions, *Thermus* enzymes were far less prone to ligate a 3'T/G mismatch. As a result, the fidelity of *Thermus* enzymes was 17- to 126-fold higher than T4 ligase (Table 2, Ligation fidelity 1). The fidelity of the newly cloned *Tsp*. AK16D ligase was similar to K294R *Tth* mutant but 6-fold higher than wild-type *Tth* enzyme. A DNA-adenylate intermediate was observed with 3'T/G mismatch

- 34 -

ligation, suggesting that a mismatch at the 3' ligation junction imposes substantial constraints on the ability of *Thermus* ligases to close the nick, thereby limiting the turnover of DNA-adenylate intermediate into ligated product and free AMP (the third step of ligation cycle). The effects of moving the T/G mismatch one base-pair away from the ligation junction was further examined. The rates of ligation with a T/G mismatch at the penultimate 3' end in general improved several-fold as compared with the T/G mismatch at the 3' end of the ligation junction. However, the ligation rates were still much slower than those of match ligation, emphasizing the importance of nucleotide complementarity near the ligation junction as well as the ultimate critical role of the perfect base-pair at the 3' end in controlling ligation reaction. Consequently, the ligation fidelity when the mismatch was at the second position from the 3' side (ligation fidelity 2) was lower than that when the mismatch was located immediately at the ligation junction. It is noteworthy that the *Tsp.* AK16D enzyme maintains extremely high fidelity (1.1×10^3) even when the mismatch is at the penultimate position, further underscoring the discriminative power of this new *Thermus* ligase.

Example 14 - Thermostable DNA Ligase Fidelity In The Presence Of Mn^{2+}

Many enzymes such as DNA polymerase and restriction endonucleases demonstrate relaxed specificity when Mn^{2+} is used as the metal cofactor. The influence of metal ion substitution on ligase fidelity has not been fully investigated although it is known that Mn^{2+} can be used as an alternative metal cofactor for a ligation reaction ((Ho, et al., *J Virol*, 71(3):1931-1937 (1997) and Cheng, et al., *Nucleic Acids Res*, 25(7):1369-1374 (1997), which are hereby incorporated by reference). The reaction rates of the match and mismatch ligation for *Tsp.* AK16D ligase and *Tth* ligase were determined. As shown in Table 3, the match ligation rates were higher with Mg^{2+} than with Mn^{2+} (Table 3), in agreement with the consistent high ligation rate under various Mg^{2+} conditions (Figure 4-6).

Table 3. DNA ligase fidelity with Mn^{2+} ^a

Ligase	Concentration (nM)	Initial rate of C-G match (fmol/min)	Initial rate of T-G mismatch (fmol/min)	Ligation fidelity
<i>Tth</i> -wt	1.25	2.6×10^1	3.7×10^{-1}	7.0×10^1
<i>Tsp.</i> AK16D	12.5	9.5×10^1	1.1×10^{-1}	8.6×10^2

5 ^a Reaction conditions were identical to those in Table 2, except that 10 mM Mn^{2+} was used in place of Mg^{2+} . Ligation fidelity was defined as the ratio of Initial Rate of C-G match divided by Initial Rate of T-G mismatch at 3'-end. Results were calculated as the average of at least two experiments.

10 The mismatch ligation rate of *Tth* ligase was about six-fold higher with Mn^{2+} than with Mg^{2+} while that of *Tsp.* AK16D ligase was about 4-fold higher. Thus, as with other previously studied DNA enzymes, DNA ligases also demonstrate relaxed specificity when Mg^{2+} is substituted with Mn^{2+} . As a result, the fidelity factors of *Tth* ligase and *Tsp.* AK16D ligase were reduced 12- and 6-fold, respectively (Tables 2-3).

15 Remarkably, the *Tsp.* AK16D enzyme retains 12-fold higher fidelity against mismatch ligation than the *Tth* enzyme. In contrast to using Mg^{2+} as the metal cofactor, *Tth* ligase did not generate DNA-adenylate intermediate during 3'T/G mismatch ligation with Mn^{2+} . This observation suggests that the nick closure of a 3'T/G mismatch by the *Tth* enzyme is accelerated with Mn^{2+} . On the other hand, the *Tsp.* AK16D enzyme

20 accumulated DNA-adenylate intermediate during 3'T/G mismatch ligation with either Mg^{2+} or Mn^{2+} . These results indicate that the nick closure of a 3'T/G mismatch with Mn^{2+} by *Tsp.* AK16D DNA ligase remains as the rate-limiting step, which accounts for the higher fidelity of this enzyme.

25 Studies on *Tth* DNA ligase has deepened understanding of thermostable ligases and has reaffirmed the common theme of ligation — adenylation of ligase at the KXDG motif (Luo, et al., Nucleic Acids Res, 24(15):3079-3085 (1996), which is hereby incorporated by reference). This study reveals that *Thermus* ligases may differ from each other as to substrate specificity despite their highly identical primary protein sequences. A highly homologous structure can be

30 anticipated from various *Thermus* ligases, but subtle local environments may dictate the probability of accepting a particular mismatch as the substrate. The fidelity of the

Thermus ligases may be determined by multiple domains, multiple motifs and/or multiple sequence elements. In comparison of *Tth* and *Tsp.* AK16D ligases, one can find that although K294R (in an identical local environment, see Figure 1B) enhances the fidelity of *Tth* ligase (Luo, et al., Nucleic Acids Res, 24(15):3071-3078 (1996), which is hereby incorporated by reference), *Tsp.* AK16D ligase with a K in this position can still demonstrate superior mismatch discrimination. Additional sequence elements remain to be uncovered. The R substitution at the adjacent position to the KXDG motif may have an effect on the *Tsp.* AK16D ligase's specificity, because studies on *Chlorella* ligase has emphasized the importance of occupying AMP binding pocket for nick recognition (Sriskanda, et al., Nucleic Acids Res, 26(2):525-531 (1998)). The accumulation of DNA-adenylate intermediate with some divalent metal ions by *Tsp.* AK16D ligase asserts that the nick closure step of a ligation reaction can be sensitive to the selection of metal ions, gapped substrates and mismatch substrates. More structural and functional studies on *Tsp.* AK16D ligase could reveal how this enzyme achieves high fidelity with different substrates and different metal ions.

Although the invention has been described in detail for the purpose of illustration, it is understood that such details are solely for that purpose and that variations can be made therein by those skilled in the art without departing from the spirit of the scope of the invention which is defined by the following claims.

- 37 -

WHAT IS CLAIMED:

1. A thermostable ligase having 100 fold higher fidelity than T4 ligase and 6 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when
5 sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent the ligation junction.
- 10 2. A thermostable ligase according to claim 1, wherein said thermostable ligase has 50 fold higher fidelity than T4 ligase and 5 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation
15 junction at the base penultimate to the ligation junction.
3. A thermostable ligase according to claim 2, wherein, in the presence of a Mn^{2+} cofactor, said thermostable ligase has a 12 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a
20 pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent to the ligation junction.
4. A thermostable ligase according to claim 3, wherein the
25 thermostable ligase has an arginine adjacent its active site lysine in the KXDG motif where X is any amino acid.
5. A thermostable ligase according to claim 1, wherein, in the presence of a Mn^{2+} cofactor, said thermostable ligase has a 12 fold higher fidelity than
30 wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a

mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent to the ligation junction.

6. A thermostable ligase according to claim 5, wherein the
5 thermostable ligase has an arginine adjacent its active site lysine in the KXDG motif where X is any amino acid.

7. A thermostable ligase according to claim 1, wherein the
10 thermostable ligase has an arginine adjacent its active site lysine in the KXDG motif where X is any amino acid.

8. A thermostable ligase according to claim 1, wherein the
15 thermostable ligase has a molecular weight of 78 to 81 kDa determined by SDS-PAGE.

9. A thermostable ligase according to claim 1, wherein the
thermostable ligase has an amino acid sequence of SEQ. ID. No. 1.

10. A thermostable ligase having 50 fold higher fidelity than T4
20 ligase and 5 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base penultimate to the ligation junction.

11. A thermostable ligase according to claim 10, wherein, in the
25 presence of a Mn^{2+} cofactor, said thermostable ligase has a 12 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation
30 junction at the base immediately adjacent to the ligation junction.

- 39 -

12. A thermostable ligase according to claim 11, wherein the thermostable ligase has an arginine adjacent its active site lysine in the KXDG motif where X is any amino acid.
- 5 13. A thermostable ligase according to claim 10, wherein the thermostable ligase has an arginine adjacent its active site lysine in the KXDG motif where X is any amino acid.
- 10 14. A thermostable ligase having an arginine adjacent its active KXDG motif where X is any amino acid.
- 15 15. A thermostable ligase having, in the presence of a Mn^{2+} cofactor, a 12 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent to the ligation junction.
- 20 16. An isolated DNA molecule encoding a thermostable ligase, wherein the thermostable ligase has a 100 fold higher fidelity than T4 ligase and 6 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent the ligation junction.
- 25 17. An isolated DNA molecule according to claim 16, wherein said thermostable ligase has 50 fold higher fidelity than T4 ligase and 5 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base penultimate to the ligation junction.
- 30

18. An isolated DNA molecule according to claim 16, wherein, in the presence of a Mn^{2+} cofactor, said thermostable ligase has a 12 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent to the ligation junction.

19. An isolated DNA molecule according to claim 16, wherein the thermostable ligase has an arginine adjacent its active site lysine in the KXDG motif where X is any amino acid.

20. An isolated DNA molecule according to claim 16, wherein said thermostable ligase has a molecular weight of 78 to 81 kDa determined by SDS-PAGE.

21. An isolated DNA molecule according to claim 16, wherein the thermostable ligase has an amino acid sequence of SEQ. ID. No. 1.

22. An isolated DNA molecule according to claim 16, wherein said DNA molecule has a nucleotide sequence of SEQ. ID. No. 2.

23. An isolated DNA molecule according to claim 16, wherein said DNA molecule hybridizes to a nucleic acid having SEQ. ID. No. 2 under stringent conditions.

24. A DNA expression system transduced with a heterologous DNA molecule according to claim 16.

25. A DNA expression system transduced with a heterologous DNA molecule according to claim 17.

26. A DNA expression system transduced with a heterologous DNA molecule according to claim 18.

27. A DNA expression system transduced with a heterologous DNA molecule according to claim 19.

28. A host cell transduced with a heterologous DNA molecule according to claim 16.

29. A host cell transduced with a heterologous DNA molecule according to claim 17.

30. A host cell transduced with a heterologous DNA molecule according to claim 18.

31. A host cell transduced with a heterologous DNA molecule according to claim 30.

32. A method for detecting, in a sample, a target nucleotide sequence which differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations, said method comprising:
providing a sample potentially containing a target nucleotide sequence which differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations;
providing one or more oligonucleotide probe sets, each characterized by (a) a first oligonucleotide probe having a target specific portion and (b) a second oligonucleotide probe having a target-specific portion, wherein the oligonucleotide probes in a particular set are suitable for hybridization to a target nucleotide sequence which differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations and for ligation together when hybridized adjacent to one another on the target nucleotide

sequence, but have a mismatch which interferes with such ligation when hybridized to any other nucleotide sequence present in the sample;

providing a thermostable ligase having 100 fold higher fidelity than T4 ligase and 6 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent the ligation junction;

blending the sample, the one or more oligonucleotide probe sets, and the thermostable ligase to form a ligase detection reaction mixture;

subjecting the ligase detection reaction mixture to one or more ligase detection reaction cycles comprising a denaturation treatment, wherein any hybridized oligonucleotides are separated from the target nucleotide sequence, and a hybridization treatment, wherein the oligonucleotide probe sets hybridize at adjacent positions in a base specific manner to their respective target nucleotide sequences, if present in the sample, and ligate to one another to form a ligation product sequence containing the target specific portions connected together with the ligation product sequences for each set being distinguishable from other nucleic acids in the ligase detection reaction mixture, wherein the oligonucleotide probe sets may hybridize to a nucleotide sequence in the sample other than their respective target nucleotide sequences but do not ligate together due to a presence of one or more mismatches and individually separate during the denaturation treatment; and

detecting the presence of ligation product sequences produced as a result of the target nucleotide sequence being present in the sample.

25

33. A method according to claim 32, wherein said thermostable ligase has 50 fold higher fidelity than T4 ligase and 5 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base penultimate to the ligation junction.

30

34. A method according to claim 33, wherein, in the presence of a Mn^{2+} cofactor, said thermostable ligase has a 12 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent to the ligation junction.

35. A method according to claim 32, wherein, in the presence of a Mn^{2+} cofactor, said thermostable ligase has a 12 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent to the ligation junction.

36. A method according to claim 32, wherein the thermostable ligase has an arginine adjacent its active site lysine in the KXDG motif where X is any amino acid.

37. A method according to claim 32, wherein the thermostable ligase has a molecular weight of 78 to 81 kDa as determined by SDS-PAGE.

38. A method according to claim 32, wherein the thermostable ligase has an amino acid sequence of SEQ. ID. No. 1.

39. A method according to claim 32 further comprising:
amplifying, prior to said blending, the target nucleotide sequence present in the sample.

40. A method according to claim 39, wherein said amplifying is carried out by polymerase chain reaction.

41. A method for detecting, in a sample, a target double stranded nucleic acid formed from first and second complementary target nucleotide sequences which differ from other nucleotide sequences by one or more single base changes, insertions, deletions, or translocations, said method comprising:

5 providing a sample potentially containing a target double stranded nucleic acid formed from first and second complementary nucleotide sequences which differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations;

providing a first oligonucleotide probe set, characterized by (a)
10 a first oligonucleotide probe having a target specific portion and (b) a second oligonucleotide probe having a target-specific portion, wherein the oligonucleotide probes in the first set are complementary to the first target nucleotide sequence which differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations and are suitable for ligation together
15 when hybridized adjacent to one another on the first target nucleotide sequence, but have a mismatch which interferes with such ligation when hybridized to any other nucleotide sequence present in the sample;

providing a second oligonucleotide probe set, characterized by
(a) a third oligonucleotide probe having a target specific portion and (b) a fourth
20 oligonucleotide probe having a target-specific portion, wherein the oligonucleotide probes in the second set are complementary to the second target nucleotide sequence which differs from other nucleotide sequences in the sample by one or more single base changes, insertions, deletions, or translocations and are suitable for ligation together when hybridized adjacent to one another on the second target nucleotide
25 sequence, but have a mismatch which interferes with such ligation when hybridized to any other nucleotide sequence present in the sample;

providing a thermostable ligase having 100 fold higher fidelity than T4 ligase and 6 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized
30 to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent the ligation junction;

- 45 -

blending the sample, the first and second oligonucleotide probe sets, and the thermostable ligase to form a ligase chain reaction mixture;

subjecting the ligase chain reaction mixture to one or more ligase chain reaction cycles comprising a denaturation treatment, wherein any
5 hybridized oligonucleotides are separated from the target nucleotide sequences, and a hybridization treatment, wherein the oligonucleotide probe sets hybridize at adjacent positions in a base specific manner to their respective target nucleotide sequences, if present in the sample, and ligate to one another to form a ligation product sequence containing the target specific portions connected together with the ligation product
10 sequences for each set being distinguishable from other nucleic acids in the ligase chain reaction mixture, wherein the oligonucleotide probe sets may hybridize to a nucleotide sequences in the sample other than their respective target nucleotide sequences but do not ligate together due to a presence of one or more mismatches and individually separate during the denaturation treatment; and
15 detecting the presence of ligation product sequences produced as a result of the target nucleotide sequence being present in the sample.

42. A method according to claim 41, wherein said thermostable ligase has 50 fold higher fidelity than T4 ligase and 5 fold higher fidelity than wild-
20 type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base penultimate to the ligation junction.

25 43. A method according to claim 42, wherein, in the presence of a Mn^{2+} cofactor, said thermostable ligase has a 12 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base
30 immediately adjacent to the ligation junction.


44. A method according to claim 41, wherein, in the presence of a Mn^{2+} cofactor, said thermostable ligase has a 12 fold higher fidelity than wild-type *Thermus thermophilus* ligase, when sealing a ligation junction between a pair of oligonucleotide probes hybridized to a target sequence where there is a mismatch with the oligonucleotide probe having its 3' end abutting the ligation junction at the base immediately adjacent to the ligation junction.

45. A method according to claim 42, wherein the thermostable ligase has an arginine adjacent its active site lysine in the KXDG motif where X is any amino acid.

46. A method according to claim 41, wherein the thermostable ligase has a molecular weight of 78 to 81 kDa as determined by SDS-PAGE.

47. A method according to claim 41, wherein the thermostable ligase has an amino acid sequence of SEQ. ID. No. 1.

1/7

**FIG. 1A**

113 YTVERKVDGLSVNLYYE 129..... 231 LEE ... TG 239..... 285 PFEADGVVVKLD 296	Tsp. AK16D
YTVEHKVDGLSVNLYYE LEE ... TG PFEADGVVVKLD	Taq. YT1
YTVEHKVDGLSVNLYYE LEEVEREG PFEADGVVVKLD	Tth
YTVEHKVDGLSVNLYYE LEEVEREG PFEADGVVVKLD	T. flavus
YTVEHKVDGLSVNLYYE LEE ... SG PFEADGVVVKMD	Tfi. Tok4A2
YTVEHKVDGLSVNLYYE LEE ... SG PFEADGVVVKLD	Tfi. Tok6A1
YTVEHKVDGLSVNLYYE LEE ... SG PFEADGVVVKLD	Tsp. SM32
YTVEHKVDGLSVNLYYE LEE ... SG PFEADGVVVKLD	Tsp. Vil3
YTVEHKVDGLSVNLYYE LEE ... SG PFEADGVVVKLD	T. scot


FIG. 1B

MTLEEARRRVNELRDLIRYHNYLYYVLDAP EISDAEYDRLLRELKELEERFPELKSPDSP	60
TEQVGARPLEATFRPVRHPTRMYSLDNAFSLDEVRAFEERIERALGRKGPFLYTVERKVD	120
GLSVNLYYEEGILVFGATRGDGETGEEVTQNLLTIPTIPRRLTGVPDRLEVRGEVYMPIE	180
AFLRLNQEELEEAGERIFKNPRNAAAGSLRQKDPRVTARRGLRATFYALGLGLEETGLKSQ	240
HDLLLWLRERGFPPVEHGFTALGAEGVEEVYQAWLKERRKLPFEADGVVVKLDDLALWRE	300
LGYTARTPRFALAYKFPAAEEKETRLLSVAFQVGRTGRITPVGVLPEVFI EGSEVSRVTLH	360
NESFIEELDVRIGDWLVHKAGGVIPEVLRVLKERRTGEEKPIIWPENCPECGHALIKEG	420
KVHRCNPPLCPAKRFEAIRHYASRKAMDIQGLGEKLEKLLKGLVRDVADLYRLKKEDL	480
VNLERMGEKSAENLLRQIEESKGRGLERLLYALGLPGVGEVLARNLALRF GHMDRLLLEAG	540
LEDLLEVEGVGELTARA ILNTLKDPEFRDLVRR LKEAGVEMEAKEREGEALKGLTFVITG	600
ELSRPREEVKALLRRLGAKVTDSVSRKTSFLVVGENPGSKLEKARALGVPTLSEEELYRL	660
IEERTGKDPRALTA	674




FIG. 1C

This Page Blank (uspto)

2/7


**FIG. 2**

This Page Blank (uspto)


**FIG. 3A****FIG. 3B****FIG. 3C**

This Page Blank (uspto)

4/7

**FIG. 4A**



LIGATION PRODUCT—

DNA ADENYLATE
INTERMEDIATES
SUBSTRATE—**FIG. 4B**

SUBSTITUTE SHEET (RULE 26)



This Page Blank (uspto)

5/7

**FIG. 5A****FIG. 5B**



This Page Blank (uspto)

6/7

**FIG. 6A****FIG. 6B**

This Page Blank (uspto)

7/7

**FIG. 7A****FIG. 7B**

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/25437

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/52 C12N9/00 C12N1/21 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, CAB Data, STRAND, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	J. TONG ET AL.: "Biochemical properties of a high fidelity DNA ligase from Thermus species AK16D" NUCLEIC ACIDS RESEARCH, vol. 27, no. 3, 1 February 1999 (1999-02-01), pages 788-794, XP002141349 IRL PRESS LIMITED, OXFORD, ENGLAND the whole document	1-31
T	M. ZIRVI ET AL.: "Ligase-based detection of mononucleotide repeat sequences" NUCLEIC ACIDS RESEARCH, vol. 27, no. 24, 15 December 1999 (1999-12-15), page e40 XP002141350 IRL PRESS LIMITED, OXFORD, ENGLAND the whole document	32-47
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

28 June 2000

Date of mailing of the international search report

14/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hornig, H

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>J. LUO AND F. BARANY: "Identification of essential residues in Thermus thermophilus DNA ligase" NUCLEIC ACIDS RESEARCH, vol. 24, no. 15, 1 August 1996 (1996-08-01), pages 3079-3085, XP002141351 IRL PRESS LIMITED, OXFORD, ENGLAND cited in the application the whole document</p>	
A	<p>J. LUO ET AL.: "Improving the fidelity of Thermus thermophilus DNA ligase" NUCLEIC ACIDS RESEARCH, vol. 24, no. 14, 1 August 1996 (1996-08-01), pages 3071-3078, XP002141352 IRL PRESS LIMITED, OXFORD, ENGLAND cited in the application the whole document</p>	
A	<p>BARANY F: "GENETIC DISEASE DETECTION AND DNA AMPLIFICATION USING CLONED THERMOSTABLE LIGASE" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, US, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, vol. 88, no. 1, 1 January 1991 (1991-01-01), pages 189-193, XP000368693 ISSN: 0027-8424 cited in the application the whole document</p>	
A	<p>JONSSON Z O ET AL: "Sequence of the DNA ligase-encoding gene from Thermus scotoductus and conserved motifs in DNA ligases" GENE, NL, ELSEVIER BIOMEDICAL PRESS. AMSTERDAM, vol. 151, no. 1, 30 December 1994 (1994-12-30), pages 177-180, XP004042633 ISSN: 0378-1119 cited in the application the whole document</p>	

-/--

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/25437

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>F. BARANY AND D.H. GELFAND: "Cloning, overexpression and nucleotide sequence of a thermostable DNA ligase-encoding gene" GENE, vol. 109, no. 1, 20 December 1991 (1991-12-20), pages 1-11, XP002141353 ELSEVIER SCIENCE PUBLISHERS, B.V., AMSTERDAM, NL; the whole document</p> <p style="text-align: center;">---</p>	
A	<p>WO 98 03673 A (CORNELL RES FOUNDATION INC ;PURDUE RESEARCH FOUNDATION (US)) 29 January 1998 (1998-01-29) cited in the application the whole document</p> <p style="text-align: center;">---</p>	
A	<p>WO 91 17239 A (CORNELL RES FOUNDATION INC ;CALIFORNIA INST OF TECHN (US)) 14 November 1991 (1991-11-14) cited in the application the whole document</p> <p style="text-align: center;">-----</p>	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/25437

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9803673	A	29-01-1998	AU	3881997 A	10-02-1998
			CA	2260818 A	29-01-1998
			EP	0956359 A	17-11-1999
WO 9117239	A	14-11-1991	EP	0528882 A	03-03-1993
			US	5494810 A	27-02-1996
			US	5830711 A	03-11-1998
			US	6054564 A	25-04-2000